

Evaluation of Oxidative Stress and Vitamin A in Patients of Chronic Otitis Media with and Without Cholesteatoma.

¹Dr. Bindiya Ghedia, ²Dr Brahma Reddy Malapati and ³Dr. Prashant D. Dave

¹Assistant professor, Department of Microbiology, Smt. B.K. Shah medical institute and Research Centre, Vadodara.

²Associate Professor, Department of Biochemistry, Dr MK Shah Medical College and Research Centre, Ahmedabad

³Associate Professor, Department of Preventive and Social Medicine, Parul Institute of Medical Science and Research, Parul University

*Corresponding Author

Dr Brahma Malapati Ready
(brahmareddy.malapati@gmail.com)

Article History

Received: 08.08.2025

Revised: 15.09.2025

Accepted: 24.10.2025

Published: 04.11.2025

Abstract:

Background: Chronic otitis media is a serious clinical condition characterized by an accumulation of fluid behind the tympanic membrane, while lacking the signs and symptoms of acute infection. The aim of this study was to investigate the effect of oxidative stress and antioxidant situation on chronic otitis media (COM) with and without cholesteatoma. **Methodology:** The study included a total of 75 cases of COM and further subcategorized based on cholesteatoma, 35 cases COM with cholesteatoma (20 females and 15 males) and 35 COM without cholesteatoma (16 females and 19 males). Serum specimens were taken prior to surgery and diseased tissue specimens from the ear were obtained during surgery from all patients. A proper audiometry testing was done. serum specimens were taken from every individual. The malondialdehyde (MDA), superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GHPx) were measured in the serum samples of the patient of COM with and without cholesteatoma. Vitamin A were determined by HPLC methodology. **Results:** The patients in this study, who might be of either gender, ranged in age from 25 to 40. Patients with chronic otitis media were subdivided based on their cholesteatoma status, as well as the levels of different oxidative indicators and vitamin A. The present study provided information on the age, gender, and BMI of each group. The present study presented data on antioxidant enzyme levels along with other biochemical parameters for both groups. **Conclusion:** The present study could help clarify the fundamental causes of the pathophysiology of COM. A finding of vitamin A does not show any direct role in occurrence of diseases but it shown significant difference in COM with and without cholesteatoma.

Keyword: Cholesteatoma; Chronic Otitis Media; Oxidative stress; vitamin A.

INTRODUCTION

The rupture of the ear drum and inflammation of the mucosa lining the middle ear hollow and the airy spaces of the temporal bone that last longer than three months are known as chronic otitis media (COM). A growing tumor made up of keratinizing squamous epithelium in the middle ear and/or mastoid process is a characteristic of COM with cholesteatoma [1]. Traditionally, the mucoperiosteum has been the exclusive site of COM pathology. Any pathology that surpasses this threshold may lead to consequences such as meningitis, osteitis, and bone deterioration [2].

Under normal physiological settings, reactive oxygen species (ROS) are significant pathogenic mediators of many illnesses. Not only are some reactive forms of oxygen necessary for human life, but they can also be harmful to the body. ROS can be harmful to many tissues because they are produced during chemical and metabolic interactions involving molecules like lipid peroxides, hydrogen peroxide, hydroxyl free radicals, and many more derivatives. Two examples of oxidative enzymes are myeloperoxidase (MPO) and nitric oxide (NO) [3].

Free oxygen radicals, or FORs, are critical for both metabolic processes and immunological responses. The

body produces FORs as defense cells such as macrophages, neutrophils, and monocytes combat antigens [4]. However, an overabundance of FORs results in tissue damage, which impedes the body's normal healing process and lengthens the duration of inflammation [5]. Free radicals are transformed into less dangerous forms by antioxidants, which also stop new free radicals from being created. After antioxidants have neutralized the ROS in a tissue, damage to the tissue, lipids, proteins, and DNA results, ultimately leading to cellular death [6]. An imbalance between increased reactive oxygen species (ROS) and antioxidant defence mechanisms interferes with the healing process of wounds and encourages persistent inflammation, which facilitates the development of cholesteatoma in COM. Antioxidant enzymes like glutathione peroxidase, catalase, and superoxide dismutase not only play a fundamental but indispensable role to fend off free radical damage via antioxidant defense system.[5]

In this study, Antioxidant enzymes, including catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GHPx), malondialdehyde (MDA), and vitamin A, were evaluated in serum samples of COM patients with and without cholesteatoma. These samples showed mucosal inflammation in the middle ear.

MATERIALS & METHODS:

A type of analytical and comparative study conducted on confirm diagnosis of chronic otitis media (COM) of total 75 patients after obtaining the ethical approval from institutional ethical committee. A confirmed cases of COM were enrolled from different ENT clinic of Cuttack and Balasore, Odisha, India. A total of 75 cases were further divided based on their cholesteatoma. 75 patients were further comprised 35 patients with COM with cholesteatoma (20 females and 15 males) and 35 COM without cholesteatoma (16 females and 19 males). Pure tone audiometry and temporal bone computed tomography (CT) were used to evaluate each patient. Participants in the study were those whose CT scans revealed soft tissue density, which indicated granulation tissue in the middle ear and mastoid cells. Antioxidant vitamin supplements, including vitamins A were not being regularly administered to any of the patients. Patients with systemic illness or acute infections were not included. Furthermore, the patients were not on any medications, nor were they smokers or drinkers. After an overnight fasting the blood samples were drawn with aseptic precaution. The serum was then separated by centrifugation at 3,000 RPM for 10 minutes. Serum samples used for the measurement of antioxidant enzymes levels such as CAT, GHPx, and SOD. For the

measurement of MDA and vitamin E, serum sample were stored at -40°C till the analysis. Every patient had a mastoidectomy. During surgery, tissue samples were extracted from problematic locations in the middle ear or mastoid cells. After splitting the specimens into two groups, one for each type of cholesteatoma the specimens were frozen at -40°C in dry tubes.

For vitamin A determination, 75 μL of serum was transferred into a 1.5 mL tube. Isopropanol (300 μL) and an internal standard (0.75 mg/L acetic retinyl) were added to the tube, and the mixture was vigorously vortexed for 2 min. Following this, the mixture was centrifuged for 15 min at 8500 revolutions per minute. The supernatant was collected, and the determination of vitamin A levels was conducted using the high-performance liquid chromatography (HPLC) method.

Statistical analysis:

Version 22.0 of the SPSS software was used to assess each data set's statistical analysis. A Pearson chi-squared test was run on the group's gender evaluation statistics. Numbers and percentages were used to report the comparison results that were obtained. The continuous variables between the groups were compared using an independent sample t-test. The results of this comparison are given as mean and standard deviation.

RESULTS:

The age ranged from 25to 40 years of the patients with either gender was enrolled in this study. A chronic otitis media patients were subgroup on the basis of their cholesteatoma condition and levels of various oxidative markers and vitamin A were determinised. Age, gender and BMI data of both the groups are presented in Table 1. Antioxidant enzymes and other Biochemical data for both the groups are presented in Table 2.

Table-1: Statistical analysis of general characteristic

Characteristic		COM with cholesteatoma (n=35)	COM without cholesteatoma (n=35)	P value
Age (Mean $\pm$ SD)		31.0 $\pm$ 11.4	35.1 $\pm$ 10.9	0.34
sex	Male	20	16	0.27
	Female	15	19	
BMI (kg/m ²)		24.8 $\pm$ 4.5	25.8 $\pm$ 4.6	0.83

Table-2: Analysis of oxidative stress and vitamin A in COM with & without cholesteatoma

Characteristic	COM with cholesteatoma (n=35)	COM without cholesteatoma (n=35)	P value
CAT (U/L)	20.55 $\pm$ 3.01	25.64 $\pm$ 5.01	<0.05
GHPx (EU/mL)	34.32 $\pm$ 3.02	34.48 $\pm$ 2.98	<0.05
SOD(EU/mL)	19.98 $\pm$ 1.98	21.65 $\pm$ 2.00	<0.05
MDA (nmol/mL)	35.24 $\pm$ 2.03	45.38 $\pm$ 2.43	<0.05
Vitamin A (ug/L)	504.66 $\pm$ 165.23	531.22 $\pm$ 170.23	<0.05

DISCUSSION:

A timeline of treatment illness course and its potential complication are the main cause for the chronic condition of otitis media. COM is still significant in the practice of otologists. The pathophysiology includes recurrent upper respiratory tract infections, immunosuppressive diseases, allergies, malnutrition, nasopharyngeal

lymphatic tissue hypertrophy, and craniofacial abnormalities. The chronicity of infection and tissue damage in COM have lately been attributed to ROS, [7]. Endogenous ROS production occurs during a variety of physiological events related to oxygen consumption. By chemically altering proteins, carbohydrates, nucleotides, and lipids, excessive ROS synthesis damages tissue and

may contribute to the etiology of a number of illnesses. The body nonenzymatic antioxidants include glutathione, tocopherole (vitamin E), ascorbic acid (vitamin C), beta-carotene (vitamin A), albumin, bilirubin, and uric acid [8]. The most significant antioxidant enzymes are SOD, GHPx, and CAT. Oxidative stress is linked to reductions in antioxidant levels or elevations in oxidant production; this state results in phospholipid peroxidation, which damages essential bodily components such lipids, lipoproteins, proteins, and DNA [9]. Polyunsaturated fatty acids are hydrolyzed into physiologically active substances at the conclusion of the process. The most significant of these substances is MDA, which signifies the body's lipid peroxidation [10]. Martanegara et al (2020) also observed that most (64.72%) cases were in the age range of 21-30 years. [11] Pratama et al conducted a study at Sanglah General Hospital, reporting that 43.2%, 18.9%, and 18.6% of CSOM cases occurred in adults, elderly, and teenagers [12]. similarly, Gaurano and Johaarjy reported that most cholesteatoma patients were aged between 20 and 35 [13] In this study, a similar trend was identified in both CSOM with and without cholesteatoma, with the highest incidence occurring in the age between 25 and 35 years. This age group appears particularly susceptible to CSOM, likely due to their productive age and potentially lower attention to hygiene, sanitation, as well as overall health.[11]

Yilmaz et al. [14] demonstrated high levels of serum MDA preoperatively, but low levels of serum SOD, GHPx, retinol, beta-carotene, alpha-tocopherole, lycopene and ascorbic acid in children with ventilation tubes due to adenoidectomy and otitis media with effusion. They showed that antioxidant levels normalized after one month postoperatively and emphasized the requirement of antioxidant therapy in these patients. Garcia Callejo et al. [15] found that the MDA value in the effusion fluid of COM patients with effusion was approximately 10 times higher than that in the middle ear fluid of OME patients. Baysal et al. [16] reported a low total antioxidant capacity and high oxidative stress index in COM patients with and without cholesteatoma.

In this study, the oxidative balance was evaluated in patients with and without cholesteatoma were compared. When the serum values of all COM patients with and without cholesteatoma were compared, then the levels of antioxidant enzyme such as MDA, SOD & CAT in the patient of COM without cholesteatoma was found to be significantly higher ($P<0.05$). The present study did not find any such differences in the mean levels of GHPx in patients of COM with and without cholesteatoma but somehow it has shown a significant difference between these two groups ($P<0.05$). These findings indicate that evaluating the values of only serum oxidative stress, excluding assessment of tissue values, may suffice for the evaluation of oxidative stress in COM patients. Also, this indicates that the severity of the disease is not

parallel to the stress oxidation, in other words, oxidative stress does not reflect the severity of the disease.

Vitamins A serve as secondary or exogenous antioxidants, effectively inhibiting and neutralizing oxidation reactions.[17] Study conducted by Baysal et al identified a significant increase in serum oxidant status and oxidative stress index among CSOM patients. Furthermore, changes in diet and nutritional intake can impact the levels of these vitamins in the blood. In this context, Vitamin A plays a role in cell differentiation and epithelial proliferation.[18] Additionally, Vitamin E, being a fat-soluble antioxidant, effectively inhibits peroxy radicals and halts the oxidation of polyunsaturated fatty acids (PUFA). According to a study by Greaves, et al, an advantage of vitamin A is its effectiveness at low concentrations of oxygen. [19] Therefore, it can complement the antioxidant properties of vitamin E that is effective at high concentrations.[20] In this study, it was observed that patients with low levels of vitamins A did not increase the risk of CSOM with cholesteatoma. Furthermore, the mean serum level of vitamin A was slightly lower in CSOM with cholesteatoma, and found significantly different ($P<0.05$). The study by Arulselvan et al stated that otitis media is a manifestation of vitamin A deficiency [20].

CONCLUSION:

Oxidative stress due to FORs plays a role in the pathogenesis of both COM with and without cholesteatoma. Assessing the values of only serum oxidative stress, excluding assessment of tissue values, may suffice for the evaluation of oxidative stress in COM patients. However, oxidative stress values may not reflect the severity of the disease. However, Vitamin A levels do not show any impact on occurrence of COM and cholesteatoma. But it has role in neutralizing the ROS, so it may have potent role in pathophysiology of diseases. For the reliable conclusion a large number of participants and details daily utility of vitamin A by patient would emphasize its role in clear outlook.

REFERENCES:

1. Semaan MT, Megerian CA. The pathophysiology of cholesteatoma. *Otolaryngol Clin North Am.* 2006 Dec;39(6):1143-59.
2. Louw L. Acquired cholesteatoma pathogenesis: stepwise explanations. *J Laryngol Otol.* 2010 Jun;124(6):587-93.
3. Sagioglu S, Ates S, Tolun FI, Oztarakci H. Evaluation of oxidative stress and antioxidants effect on turning process acute otitis media to chronic otitis media with effusion. *Niger J Clin Pract.* 2019;22(3):375-379. doi:10.4103/njcp.njcp_126_18.
4. Yilmaz T, Kocan EG, Besler HT, Yilmaz G, Gursel B. The role of oxidants and antioxidants in otitis media with effusion in children. *Otolaryngol Head Neck Surg.* 2004 Dec;131(6):797-803.

5. Shukla GK, Mahajan A, Pandey S, Gujrati VR, Vrat S, Mishra SC, et al. A study of free radicals and scavenging enzyme in tonsillitis. *Boll Chim Farm.* 1996 Dec;135(11):653-5
6. Sagioglu, S; Ates, S1; Tolun, FI2; Oztarakci, H3. Evaluation of Oxidative Stress and Antioxidants Effect on Turning Process Acute Otitis Media to Chronic Otitis Media with Effusion. *Nigerian Journal of Clinical Practice* 22(3):p 375-379, March 2019. | DOI: 10.4103/njcp.njcp_126_18
7. Baysal E, Aksoy N, Kara F, Taysi S, Taskin A, Bilinc H, et al. Oxidative stress in chronic otitis media. *Eur Arch Otorhinolaryngol.* 2013 Mar; 270(4):1203-8. 12. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007;39(1):44-84.
8. Shukla MK, Mahajan A, Pandey S, Gujrati VR, Vrat S, Mishra SC, et al. A study of free radicals and scavenging enzyme in tonsillitis. *Boll Chim Farm.* 1995 Dec;135(11):656.
9. Yoshioka T, Kawada K, Shimada T, Mori M. Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood. *Am J Obstet Gynecol.* 1979 Oct;135(3): 372-6
10. Martanegara IF, Purwanto B, Boesoirie SF. Characteristic of intratemporal complication in chronic suppurative otitis media patient with and without cholesteatoma at hasan sadikin general hospital bandung. *J Med Heal.* 2020;2(6):197–204.
11. Arya IP, Pratama A, Sudipta IM, Andi K, Saputra D. Description of patients with chronic suppurative otitis media at sanglah hospital, Denpasar, 2014–2016. *E Jurnal Med.* 2019;8(4):1
12. Gaurano JL, Joharjy IA. Middle ear cholesteatoma: characteristic CT findings in 64 patients. *Ann Saudi Med.* 2004;24(6):442–447. doi:10.5144/ 0256-4947.2004.442 2
13. Yariktas M, Doner F, Dogru H, Yasan H, Delibas N. The role of free oxygen radicals on the development of otitis media with effusion. *Int J Pediatr Otorhinolaryngol.* 2004 Jul;68(7):889-94.
14. Garcia Callejo FJ, Estors Ferrero J, Morant Ventura A, Segarra Cortes P, Velert Vila MM. Lipoperoxidation in otorrhea of the middle ear as a marker of infection: clinical application. *Acta Otorrinolaringol Esp.* 2000 Aug-Sep;51(6):478-81
15. Baysal E, Aksoy N, Kara F, Taysi S, Taskin A, Bilinc H, et al. Oxidative stress in chronic otitis media. *Eur Arch Otorhinolaryngol.* 2013 Mar; 270(4):1203-8
16. Lee GY, Han SN. The role of vitamin E in immunity. *Nutrients.* 2018;10(11):1–18. doi:10.3390/nu10111614
17. Schmitz J, West KP, Khatry SK, et al. Vitamin A supplementation in preschool children and risk of hearing loss as adolescents and young adults in rural Nepal: randomised trial cohort follow-up study. *BMJ.* 2012;344(7841):1–12. doi:10.1136/bmj.d7962
18. Greaves RF, Woollard GA, Hoad KE, et al. Laboratory medicine best practice guideline: vitamins A, E and the carotenoids in blood. *Clin Biochem Rev.* 2014;35(2):81–113. doi:10.1186/bcr2338
19. Arulselvan P, Fard MT, Tan WS, et al. Role of antioxidants and natural products in inflammation. *Oxid Med Cell Longev.* 2016;2016. doi:10.1155/2016/5276130
20. Mehmet Fatih Garça, Mahfuz Turan, Barış Avşar, Ferhat Kalkan2 ·Halit Demir, Ahmet Kozan, Nazım Bozan. The Evaluation of Oxidative Stress in the Serum and Tissue Specimens of Patients With Chronic Otitis Media. *Clinical and Experimental Otorhinolaryngology.* 2015; 8 (2): 97-101.